
EXHIBIT 3



UNIVERSITY OF CALIFORNIA Agriculture

and Natural Resources

http://anrcatalog.ucdavis.edu

Lawn Watering Guide for California

Janet Hartin is UC Cooperative Extension Farm Advisor in Environmental Horticulture for San Bernardino County; Pamela M. Geisel is UCCE Farm Advisor in Environmental Horticulture for Fresno County; and Carolyn L. Unruh is a UCCE staff writer for Fresno County. This publication is adapted from "Dry Wit," a 1993 informational brochure published by the University of California, Division of Agriculture and Natural Resources.

The techniques described in this publication will help homeowners set up timed irrigation controllers for home lawns. The simple procedure involves identifying your home's climatic region, the type of turfgrass you have, and the output of your irrigation system. A set of tables provides a general guideline for scheduling lawn irrigation based on average weather data and turfgrass growth characteristics. Environmental conditions will vary somewhat from year to year and from location to location within a region, so your irrigation controller will continue to need minor adjustments from time to time in order to deliver optimum results. Ideally, you should plan to re-set your automatic irrigation system monthly during the growing seasons in response to changes in the weather.

HOW TO USE THE LAWN WATERING GUIDE

Step 1. Determine what type of lawn you have.

- Warm-season grasses include hybrid bermudagrass, common bermudagrass, zoysiagrass, St. Augustinegrass, and kikuyugrass. Dichondra is a broadleaf groundcover with water requirements similar to those of warm-season turfgrass.
- Cool-season grasses include tall fescue, Kentucky bluegrass, annual and perennial ryegrass, and bentgrass.

Step 2. Determine the output of your sprinklers.

- Set out six or more straight-sided containers of the same type, spaced evenly on your lawn. Empty tuna cans, cat food cans, or coffee mugs work well. Run the sprinklers for 20 minutes and use a ruler to measure (in inches) the depth of water in each can. To determine the average depth of water applied to the lawn, total the water depths for all of the containers and divide the total amount by the number of containers you used.
- Multiply the average depth by three to determine how many inches of water your sprinkler system applies per hour.



Sample calculation:

Can #1	1/2 inch (0.500 inch)
Can #2	5/8 inch (0.625 inch)
Can #3	1/2 inch (0.500 inch)
Can #4	3/8 inch (0.375 inch)
Can #5	1/2 inch (0.500 inch)
Can #6	3/8 inch (0.375 inch)
OR 6 CANS	2 7/8 inches (2.875 inches

TOTAL FOR 6 CANS 2 7/8 inches (2.875 inches)

2.875 inches = 0.479 inch/can (average depth per can)


6 cans

0.479 inch x 3 = 1.437 inches (nearly 1 1/2 inches) per hour

Step 3. Determine how many minutes you need to water your lawn each week.

- Locate your geographic region on the map in this publication and then find the table that matches that region. Identify your predominant lawn type (warm-season or cool-season) and the current month on the table, and match that information to the sprinkler output that you calculated in Step 2. The resulting value is the total number of minutes that you should run your sprinklers per week to provide adequate irrigation for your lawn during the current month.
- Deeper, less frequent irrigations are best for most lawns as they promote deep root growth. If you notice excessive runoff or brown spots in the lawn with once-a-week watering, however, or if your controller can't accommodate such large blocks of time, divide the total minutes per week by 2, 3, or 4 and spread the weekly water application over the week. Desert areas, slopes, or areas with shallow soils usually need several shorter watering times instead of a single weekly irrigation. Cool-season grasses have shallow roots, and often require more frequent irrigations that warm-season grasses.
- If brown spots continue to occur despite more frequent irrigations, you may
 need to check your sprinkler system's coverage for uniformity. Look for and
 replace broken, clogged, or malfunctioning sprinkler heads that deprive lawn
 areas of sufficient water. After you upgrade, repair, or significantly change your
 system, your will need to repeat Steps 2 and 3 to make sure you're applying the
 right amount of water each week.
- During periods of unseasonably high rainfall, you should shut your irrigation system off temporarily to conserve groundwater and take advantage of the unexpected moisture. Be sure to restart your system after the rainy weather has passed. Extremely hot, dry, or windy conditions may require extra irrigation to compensate for excessive water loss from the lawn.

MAP OF CALIFORNIA



Region 8: Central California Coast

Region 9: Southern California Coast

Region 10: Southern California Inland Valleys

Region 11: Southern California Deserts

Region 1: Northern California Coast

Warm-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

Cool-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

	0.5 in	1.0 in	1.5 in	2.0 in	110	0.5 in	1.0 in	1.5 in	2.0 in
JAN					JAN	15	07	05	04
FEB					FEB	36	18	12	09
MAR					MAR	55	27	18	14
APR	W	arm-seasc	n		APR	67	34	22	17
MAY	tur	fgrasses a	ire		MAY	88	44	29	22
UN	not	recommen	ded		JUN	97	48	32	24
UL	in	this region	n.		JUL	95	47	32	24
NUG					AUG	90	45	30	23
EP					SEP	76	38	25	19
OCT					OCT	48	24	16	12
VOV					NOV	32	16	11	08
DEC					DEC	21	11	07	05

Region 2: Northern Inland Valleys

Warm-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

Cool-Season Turfgrasses

Minutes per week to irrigate i	f
your hourly sprinkler output is	5:

	0.5 in	1.0 in	1.5 in	2.0 in		0.5 in	1.0 in	1.5 in	2.0 in
JAN	19	09	06	05	JAN	25	13	08	06
FEB	32	16	11	08	FEB	42	21	14	11
MAR	50	25	17	13	MAR	67	34	22	17
APR	69	35	23	17	APR	92	46	31	23
MAY	101	50	34	25	MAY	134	67	45	34
JUN	126	63	42	32	JUN	168	84	56	42
JUL	132	66	44	33	JUL	176	88	59	44
AUG	120	60	40	30	AUG	160	80	53	40
SEP	95	47	32	24	SEP	126	63	42	32
OCT	57	28	19	14	OCT	76	38	25	19
NOV	25	13	08	06	NOV	34	17	11	08
DEC	13	06	04	03	DEC	17	08	06	04

Region 3: Northeastern California Mountain Valleys

Warm-Season	Turfgrasses
-------------	-------------

Minutes per week to irrigate if your hourly sprinkler output is:

Cool-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

	0.5 in 1.0 in 1.5 in	2.0 in		0.5 in	1.0 in	1.5 in	2.0 in	
JAN			JAN	17	08	06	04	
FEB			FEB	34	17	11	08	
MAR			MAR	59	29	20	15	
APR	Warm-season		APR	101	50	34	25	
MAY	turfgrasses are		MAY	134	67	45	34	
JUN	not recommended		JUN	168	84	56	42	
JUL	in this region.		JUL	210	105	70	53	
AUG	904 134400		AUG	176	88	59	44	
SEP			SEP	126	63	42	32	
OCT			OCT	76	38	25	19	
NOV			NOV	25	13	09	06	
DEC			DEC	17	09	06	04	

Region 4: Sacramento Valley

Warm-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

Cool-Season Turfgrasses

Minutes	per week to in	rigate if
your hou	urly sprinkler o	utput is:

,					, ,					
	0.5 in	1.0 in	1.5 in	2.0 in		0.5 in	1.0 in	1.5 in	2.0 in	
IAN	19	09	06	05	JAN	25	13	08	06	
FEB	44	22	15	11	FEB	59	29	20	15	
MAR	69	35	23	17	MAR	92	46	31	23	
APR	101	50	34	25	APR	134	67	45	34	
MAY	126	63	42	32	MAY	168	84	56	42	
UN	158	79	53	39	JUN	210	105	70	53	
UL	164	82	55	41	JUL	218	109	73	55	
AUG	145	72	48	36	AUG	193	97	64	48	
SEP	113	57	38	28	SEP	151	76	50	38	
OCT	82	41	27	20	OCT	109	55	36	27	
NOV	38	19	13	09	NOV	50	25	17	13	
DEC	19	09	06	05	DEC	25	13	08	06	

Region 5: San Joaquiin Valley

Warm-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

Cool-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

,					,			•		
	0.5 in	1.0 in	1.5 in	2.0 in		0.5 in	1.0 in	1.5 in	2.0 in	
JAN	19	09	06	05	JAN	25	13	08	06	
FEB	38	19	13	09	FEB	50	25	17	13	
MAR	69	35	23	17	MAR	92	46	31	23	
APR	101	50	34	25	APR	134	67	45	34	
MAY	132	66	44	33	MAY	176	88	59	44	
JUN	164	82	55	41	JUN	218	109	73	55	
JUL	170	85	57	43	JUL	227	113	76	57	
AUG	145	72	48	36	AUG	193	97	64	48	
SEP	113	57	38	28	SEP	151	76	50	38	
OCT	69	35	23	17	OCT	92	46	31	23	
NOV	32	16	11	80	NOV	42	21	14	11	
DEC	13	06	04	03	DEC	17	08	06	04	

Region 6: Central Inland Valleys

Warm-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

Cool-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

	0.5 in	1.0 in	1.5 in	2.0 in		0.5 in	1.0 in	1.5 in	2.0 in	
JAN	32	16	11	08	JAN	42	21	14	11	
FEB	44	22	15	11	FEB	59	29	20	15	
MAR	69	35	23	17	MAR	92	46	31	23	
APR	95	47	32	24	APR	126	63	42	32	
MAY	113	57	38	28	MAY	151	76	50	38	
JUN	113	57	38	28	JUN	151	76	50	38	
JUL	132	66	44	33	JUL	176	88	59	44	
AUG	126	63	42	32	AUG	168	84	56	42	
SEP	107	54	36	27	SEP	143	71	48	36	
OCT	76	38	25	19	OCT	101	50	34	25	
NOV	44	22	15	11	NOV	59	29	20	15	
DEC	32	16	11	08	DEC	42	21	14	11	

Region 7: Sierra

Warm	-Season Tur	fgras	ses		Cool-Season Turfgrasses						
	es per weel nourly sprin				Minutes per week to irrigate if your hourly sprinkler output is:						
	0.5 in 1.	0 in	1.5 in	2.0 in		0.5 in	1.0 in	1.5 in	2.0 in		
JAN					JAN	31	15	10	8		
FEB					FEB	43	22	14	11		
MAR					MAR	79	39	26	20		
APR	Warm	-seaso	n		APR	124	62	41	31		
MAY	turfgra	isses a	re		MAY	164	82	55	41		
JUN	not reco	mmen	ded		JUN	207	103	69	52		
JUL	in this	region	١.		JUL	231	115	77	58		
AUG		_			AUG	198	99	66	50		
SEP					SEP	141	70	47	35		
OCT					OCT	96	48	32	24		
NOV					NOV	40	20	13	10		
DEC					DEC	20	10	7	5		

Region 8: Central California Coast

Warm	-Season	Turfgras	sses		Cool-Season Turfgrasses						
Minutes per week to irrigate if your hourly sprinkler output is:					Minutes per week to irrigate if your hourly sprinkler output is:						
	0.5 in	1.0 in	1.5 in	2.0 in	-	0.5 in 1.0 in		1.5 in	2.0 in		
JAN	38	19	13	09	JAN	50	25	17	13		
FEB	50	25	17	13	FEB	67	34	22	17		
MAR	63	32	21	16	MAR	84	42	28	21		
APR	88	44	29	22	APR	118	59	39	29		
MAY	101	50	34	25	MAY	134	67	45	34		
JUN	113	57	38	28	JUN	151	76	50	38		
JUL	95	47	32	24	JUL	126	63	42	32		
AUG	113	57	38	28	AUG	151	76	50	38		
SEP	95	47	32	24	SEP	126	63	42	32		
OCT	69	35	23	17	OCT	92	46	31	23		
NOV	50	25	17	13	NOV	67	34	22	17		
DEC	38	19	13	09	DEC	50	25	17	13		

Region 9: Southern California Coast

Warm-Se	ason	Turfarasses
---------	------	--------------------

Minutes per week to irrigate if your hourly sprinkler output is:

Cool-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

			-							
3 200	0.5 in	1.0 in	1.5 in	2.0 in		0.5 in	1.0 in	1.5 in	2.0 in	
JAN	44	22	15	11	JAN	59	29	20	15	
FEB	57	28	19	14	FEB	76	38	25	19	
MAR	63	32	21	16	MAR	84	42	28	21	
APR	76	38	25	19	APR	101	50	34	25	
MAY	88	44	29	22	MAY	118	59	39	29	
JUN	95	47	32	24	JUN	126	63	42	32	
JUL	107	54	36	27	JUL	143	71	48	36	
AUG	95	47	33	24	AUG	126	63	42	32	
SEP	82	41	27	20	SEP	109	55	36	27	
OCT	69	35	23	17	OCT	92	46	31	23	
NOV	50	25	17	13	NOV	67	34	22	17	
DEC	38	19	13	9	DEC	50	25	17	13	

Region 10: Southern California Inland Valleys

Warm-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

Cool-Season Turfgrasses

Minutes per week to irrigate if your hourly sprinkler output is:

	0.5 in	1.0 in	1.5 in	2.0 in		0.5 in	1.0 in	1.5 in	2.0 in
JAN	42	21	14	10	JAN	56	28	19	14
FEB	57	28	19	14	FEB	75	38	25	19
MAR	80	40	27	20	MAR	106	53	35	27
APR	96	48	32	24	APR	128	64	43	32
MAY	119	60	40	29	MAY	159	80	53	40
JUN	144	72	48	36	JUN	193	96	64	48
JUL	165	83	55	41	JUL	221	110	74	55
AUG	155	77	52	39	AUG	207	103	69	52
SEP	124	62	41	31	SEP	165	82	55	41
OCT	88	44	29	22	OCT	117	59	39	29
NOV	54	27	18	14	NOV	73	36	24	18
DEC	42	21	14	10	DEC	55	28	19	14

Region 11: Southern California Deserts

Warm-Season Turfgrasses

asses Cool-Season Turfgrasses irrigate if Minutes per week to irrigate if

Minutes per	week to	irrigate if
your hourly	sprinkler	output is:

your hourly sprinkler output is:				your hourly sprinkler output is:						
	0.5 in	1.0 in	1.5 in	2.0 in		0.5 in	1.0 in	1.5 in	2.0 in	
JAN	54	27	18	14	JAN	65	32	22	17	
FEB	75	38	25	19	FEB	90	46	30	23	
MAR	121	61	40	30	MAR	145	73	48	36	
APR	165	83	55	41	APR	198	100	66	49	
MAY	211	106	70	53	MAY	253	127	84	64	
JUN	243	121	81	61	JUN	292	145	97	73	
JUL	251	126	84	63	JUL	301	151	101	76	
AUG	218	109	73	54	AUG	262	131	88	65	
SEP	180	90	60	45	SEP	216	108	72	54	
OCT	121	61	40	30	OCT	145	73	48	36	
NOV	69	35	23	17	NOV	83	42	28	20	
DEC	43	22	14	11	DEC	52	26	17	13	

REFERENCES

- Anonymous. 2000. Best management practices to reduce production of organic materials in landscape plantings. Oakland: University of California Division of Agriculture and Natural Resources and State of California Integrated Waste Management Board.
- California Agricultural Technology Institute, Center for Irrigation Technology, online at http://www.wateright.org/
- Hartin, J., and M. J. Henry. 1999. Reusing turfgrass clippings to improve turfgrass health and performance in southern California. University of California Cooperative Extension/California Integrated Waste Management Board leaflet.
- Hartin, J., and A. Harivandi. 1999. Reusing turfgrass clippings to improve turfgrass health and performance in central and northern California. University of California Cooperative Extension/California Integrated Waste Management Board leaflet.

FOR MORE INFORMATION

You'll find detailed information on many aspects of vegetable production in these titles and in other publications, slide sets, and videos from UC ANR:

UC IPM Turfgrass Pest Management Guidelines, publication 3365-T

Evaluating Turfgrass Sprinkler Irrigation Systems, publication 21503

Healthy Turf: First Line of Defense Against Weeds, video v94-P

To order these products, visit our online catalog at http://anrcatalog.ucdavis.edu. You can also place orders by mail, phone, or fax, or request a printed catalog of publications, slide sets, and videos from

University of California Agriculture and Natural Resources Communication Services 6701 San Pablo Avenue, 2nd Floor Oakland, CA 94608-1239

Telephone: (800) 994-8849 or (510) 642-2431, FAX: (510) 643-5470

FAX: (510) 643-5470

E-mail inquiries: danrcs@ucdavis.edu

An electronic version of this publication is available on the ANR Communication Services website at http://anrcatalog.ucdavis.edu.

Publication 8044

Funding for this publication was made possible threough a grant form the Elvenia J. Slosson Fund,

©2001 by the Regents of the University of California, Division of Agriculture and Natural Resources. All rights reserved.

The University of California prohibits discrimination against or harassment of any person employed by or seeking employment with the University on the basis of race, color, national origin, religion, sex, physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (special disabled veteran, Vietnam-era veteran or any other veteran who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized). University Policy is intended to be consistent with the provisions of applicable State and Federal laws.

Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6th floor, Oakland, CA 94612-3550; (510) 987-0096. For a free catalog of other publications, telephone (800) 994-8849.

pr10/01-WJC/VFG

ISBN 978-1-60107-222-1



This publication has been anonymously peer reviewed for technical accuracy by University of California scientists and other qualified professionals. This review process was managed by the ANR Associate Editor for Environmental Horticulture.